



## King's Research Portal

*Document Version*  
Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Matcham, F., Galloway, J., Hotopf, M., Roberts, E., Scott, I. C., Steer, S., & Norton, S. (2019). Letter to the editor: The impact of targeted rheumatoid arthritis pharmacologic treatment on mental health: A systematic review and meta-analysis: Reply. *Arthritis and Rheumatology*.

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# **Letter to the editor: The impact of targeted rheumatoid arthritis pharmacologic treatment on mental health: A systematic review and meta-analysis: Reply**

Matcham F<sup>1</sup>, Galloway J<sup>2,3</sup>, Hotopf M<sup>1</sup>, Roberts E<sup>1,4</sup>, Scott IC<sup>5</sup>, Steer S<sup>3</sup>, Norton S<sup>2,6</sup>.

1. Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK.
2. Academic Rheumatology, Faculty of Life Sciences and Medicine, King's College London, UK.
3. Rheumatology Department, King's College Hospital NHS Foundation Trust, London, UK.
4. National Addication Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK.
5. Primary Care and Health Sciences, Keele University, UK.
6. Health Psychology, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK.

Corresponding author:

Dr Faith Matcham, Department of Psychological Medicine, 3<sup>rd</sup> Floor East Wing, Institute of Psychiatry, Psychology & Neuroscience, 16 De Crespigny Park, London SE5 8AF.

Telephone: 0207 848 0868

Email: [faith.matcham@kcl.ac.uk](mailto:faith.matcham@kcl.ac.uk)

Word count: 416.

To the Editor,

We thank Dr Zembrzuska for her letter relating to our systematic review and network meta-analysis describing the impact of targeted pharmacological treatments on mental health outcomes in rheumatoid arthritis (RA) [1]. She highlights clearly the disparity in approach towards managing physical versus mental comorbidities in rheumatological practice. Approximately one in four people with RA will also have symptoms of depression [2] and there has been extensive recent research conducted by us here at King's College London, and by colleagues worldwide emphasising the impact common mental disorders such as symptoms of depression and anxiety may have on long-term physical health outcomes and treatment response in RA [3-6]. The consequences of mental comorbidity in the context of RA are clear: patients have significantly impaired long-term functioning and may not respond well to expensive treatments.

There are some small-scale local examples of excellent progress made in this area. The Integrating Mental and Physical Healthcare: Research, Training and Services (IMPARTS) project provides a generic platform for web-based patient reported outcome measurement including mental health screening which allows all patients attending general hospital outpatient appointments to be assessed for symptoms of depression and anxiety as part of routine clinical practice [7]. This system links with electronic health records, allowing routinely collected physical and psychological symptoms to be immediately available for review by the treating clinician, with advice for onward referrals and links with hospital and community-based mental health services where available. In the UK, the NHS-funded programme Improving Access to Psychological Therapies (IAPT) began offering new services for people with long term conditions in 2018, which may go some way to address this unmet need. Whilst there are also some examples of dedicated psychology time allocated to rheumatology clinics, there is a huge need for standardisation in the funding of such roles to ensure all patients have equal access to psychosocial support.

Depression and anxiety in the context of physical disease are treatable [8-9] and data from other disease areas suggests that successfully managing psychiatric co-morbidities is likely to be cost-effective [10-11]. The implementation of effective mental health care into routine practice is dependent on commissioning policies funding sustainable systems of integrated health management, changes in organisational structure and professional flexibility, and the design of pragmatic interventions relevant to the context of delivery. Providing dedicated mental health care within the context of clinical disease management, as suggested by Dr Zembrzuska, is essential to achieve parity of esteem and improve patients' physical health, response to treatment, and overall quality-of-life.

## References

1. Matcham F, Galloway J, Hotopf M, *et al.* The impact of targeted rheumatoid arthritis pharmacologic treatment on mental health. *Arthritis Rheum* 2018;**70**:377-91. doi: 10.1002/art.40565.
2. Matcham F, Rayner L, Steer S, *et al.* The prevalence of depression in rheumatoid arthritis: a systematic review and meta-analysis. *Rheumatology (Oxford)* 2013;**52**:2136-48. doi: 10.1093/rheumatology/ket169.
3. Rathbun AM, Harrold LR, Reed GW. Temporal effect of depressive symptoms on the longitudinal evolution of rheumatoid arthritis disease activity. *Arthritis Care Res* 2015;**67**:765-75. doi:10.1002/acr.22515.
4. Michelsen B, Kristianslund K, Sexton J, *et al.* Do depression and anxiety reduce the likelihood of remission in rheumatoid arthritis and psoriatic arthritis? Data from the prospective multicentre NOR-DMARD study. *Ann Rheum Dis* 2017;**76**:1906-10. doi: 10.1136/annrheumdis-2017-211284.
5. Matcham F, Davies R, Hotopf M, *et al.* The relationship between depression and biologic treatment response in rheumatoid arthritis: An analysis of the British Society for Rheumatology Biologics Register. *Rheumatology (Oxford)* 2018; **57**:835-43. doi: 10.1093/rheumatology/kex528.
6. Matcham F, Norton S, Scott DL, *et al.* Symptoms of depression and anxiety predict treatment response and long-term physical health outcomes in rheumatoid arthritis: Secondary analysis of a randomized controlled trial. *Rheumatology (Oxford)* 2016;**55**:268-78. doi: 10.1093/rheumatology/kev306.
7. Rayner L, Matcham F, Hutton J *et al.* Embedding integrated mental health assessment and management in general hospital settings: Feasibility, acceptability and the prevalence of common mental disorder. *Gen Hosp Psychiatr* 2014;**36**:318-24. doi: 10.1016/j.genhosppsy.2013.12.004.
8. Rayner L, Price A, Evans A, *et al.* Antidepressants for depression in physically ill people. *Cochrane Database Syst. Rev.* 2010;:CD007503. doi:10.1002/14651858.CD007503.pub2.
9. Matcham F, Rayner L, Hutton J *et al.* Self-help interventions for symptoms of depression, anxiety and psychological distress in patients with physical illnesses: A systematic review and meta-analysis. *Clin Psychol Rev* 2014;**34**:141-57. doi: 10.1016/j.cpr.2014.01.005.
10. Walker S, Walker J, Richardson G, *et al.* Cost-effectiveness of combining systematic identification and treatment of co-morbid major depression for people with chronic diseases: the example of cancer. *Psychol Med* 2014;**44**:1451-60. doi: 10.1017/S0033291713002079.
11. Katon W, Russo J, Lin EHB, *et al.* Cost-effectiveness of a multicondition collaborative care intervention: a randomized controlled trial. *Arch Gen Psychiatry* 2012;**69**:506-14. doi: 10.1001/archgenpsychiatry.2011.1548.